

(12) **EUROPEAN PATENT APPLICATION**

(21) Application number: 86306325.1

(22) Date of filing: 15.08.86

(51) Int. Cl.⁴: **C07D 285/12**, **C07D 285/16**,
C07D 271/10, **C07D 253/06**,
C07D 417/04, **C07K 5/06**,
A61K 31/41, **A61K 31/53**,
A61K 31/535, **A61K 31/54**,
A61K 37/02

(30) Priority: 31.08.85 GB 8521697
09.10.85 GB 8524932
06.12.85 GB 8530143
22.04.86 GB 8609793
17.05.86 GB 8612060

(43) Date of publication of application:
08.04.87 Bulletin 87/15

(84) Designated Contracting States:
AT DE NL SE

(71) Applicant: **FISONS plc**
Fison House Princes Street
Ipswich Suffolk IP1 1QH(GB)


(72) Inventor: **Brown, Roger Charles**
8 Duncan Way Gorse Covert
Loughborough Leicestershire(GB)
Inventor: **Robinson, David Hulme**
12 Grassholme Drive
Loughborough Leicestershire LE11 0NS(GB)
Inventor: **Dixon, John**
Church Farmhouse Main Street Great Dalby
Melton Mowbray Leicestershire(GB)

(74) Representative: **Craig, Christopher Bradberry**
et al
Fisons plc 12 Derby Road
Loughborough Leicestershire LE11 0BB(GB)

(54) **5- and 6- Membered heterocyclic ring angiotensin converting enzyme inhibitors.**

EP 0 217 519 A1

(57) There are described compounds of formula I,



I

in which Y is S, O or NR_n,

n is 0 or 1,

R₁ is hydrogen or alkyl C 1 to 10,

R₂ is hydrogen, alkyl C 1 to 10, cycloalkyl C3 to 10, CF₃, SR₁₀, a 5 or 6 membered heterocyclic group containing one or more S, O or N atoms, NR₄R₅, phenyl or phenylalkyl C7 to 12, the phenyl, phenylalkyl and heterocyclic groups optionally being fused to a further phenyl group, the heterocyclic group and any phenyl group optionally being substituted by alkyl C 1 to 6, halogen, alkoxy C 1 to 6, nitro, nitrile, CF₃, SR₄, NR₂R₁₁, or hydroxy,

R₄, R₅ and R₁₁, which may be the same or different, are each hydrogen or alkyl C 1 to 10,

R₆ and R₈, which may be the same or different, are each hydrogen, alkyl C 1 to 10 or phenyl,

R₁₀ is alkyl C 1 to 10,

X₁ is S or O, and

D is a chain comprising from 2 -16 atoms, which chain carries an O or S containing substituent at a position 2 -6 atoms away from the group C=X₁,

and pharmaceutically acceptable salts, esters and amides thereof,


there are also described methods for making the compounds and pharmaceutical formulations, eg for the treatment of hypertension, containing them.

This invention relates to new compounds, methods for their preparation and compositions containing them.

A wide variety of angiotensin converting enzyme (ACE) inhibitors are known, eg from French Patent Specification No. 2,372,804 and European Patent Specification No. 0012401.

We have now found a group of compounds having advantageous properties, eg as ACE inhibitors.

According to the invention we provide compounds of formula I,



I

in which Y is S, O or NR₁,

n is 0 or 1,

R₁ is hydrogen or alkyl C 1 to 10,

R₂ is hydrogen, alkyl C 1 to 10, cycloalkyl C3 to 10, CF₃, SR₁₀, a 5 or 6 membered heterocyclic group containing one or more S, O or N atoms, NR₁R₂, phenyl or phenylalkyl C7 to 12, the phenyl, phenylalkyl and heterocyclic groups optionally being fused to a further phenyl group, the heterocyclic group and any phenyl group optionally being substituted by alkyl C 1 to 6, halogen, alkoxy C 1 to 6, nitro, nitrile, CF₃, SR₁₀, NR₁R₂, or hydroxy,

R₃, R₄ and R₁₁, which may be the same or different, are each hydrogen or alkyl C 1 to 10,

R₅ and R₆, which may be the same or different, are

each hydrogen, alkyl C 1 to 10 or phenyl,

R₁₀ is alkyl C 1 to 10,

X₁ is S or O, and


D is a chain comprising from 2 -16 atoms, which chain carries an O or S containing substituent at a position 2 -6 atoms away from the group C=X₁,

and pharmaceutically acceptable salts, esters and amides thereof.

According to the invention we also provide a process for the production of a compound of formula I, or a pharmaceutically acceptable salt, ester or amide thereof, which comprises

a) removal of a protecting group from a compound of formula I in which one or more of the amino or carboxylic acid groups is protected,

b) reaction of a compound of formula II,




II

55

or a salt, ester, amide, tautomer, or protected derivative thereof,

in which R_3 , Y and n are as defined above,

with a compound of formula III,




in which D and X, are as defined above, and

X is a good leaving group

c) conversion of a compound of formula I in which the asymmetric carbon atom of the Y containing heterocyclic ring is in the R configuration into a corresponding compound in which that carbon atom is in the S configuration,

d) reaction of a compound of formula II, in which R_3 , Y and n are as defined above,

with a compound of formula VI,



in which D and X, are as defined above, or

e) production of a pharmaceutically acceptable salt of a compound of formula I, by treating a compound of formula I, or another salt, an ester or an amide thereof, with a compound containing an available pharmaceutically acceptable ion and capable of converting the compound of formula I or the other salt, ester or amide thereof, to a pharmaceutically acceptable salt of the compound of formula I,

and where desired or necessary deprotecting the resulting compound, or converting a compound of formula I to a pharmaceutically acceptable salt, ester or amide thereof or vice versa.

In process a) the protecting group can be any convenient protecting group conventionally used in peptide synthesis and may be removed using techniques conventionally used in peptide synthesis. Thus carboxy protecting groups which may be used are alkoxy C 1 to 6, which may be a straight chain or branched alkoxy, eg t-butyloxy; or phenylalkoxy C7 to 12, eg benzyloxy. These groups can be removed by hydrolysis, for example basic hydrolysis, eg using aqueous methanolic sodium hydroxide; or cleavage using, for example, trifluoroacetic acid; or by hydrogenation, eg using palladium on charcoal. Amino-protecting groups which may be mentioned include alkylloxycarbonyl C2 to 7, eg t-butyloxycarbonyl or phenylalkyloxycarbonyl C8 to 13, eg benzyloxycarbonyl. We prefer to use starting materials in which the carboxy groups are protected.

In process b) the group X may be halo, eg bromo or chloro. The reaction may be carried out in a solvent which is inert under the reaction conditions, eg acetonitrile, at a temperature of from 0° to 100°C, preferably at about 30°C. The reaction is preferably carried out under basic conditions, eg in the presence of triethylamine or polyvinylpyridine.

The reaction of process c) may be carried out in a solvent which is inert under the reaction conditions, eg acetonitrile, at a temperature of from 0°C to the boiling point of the solvent, preferably of from 20° to 30°C. The reaction may be carried out under anhydrous conditions, eg in the presence of molecular sieves, and in the presence of a base, eg pyrrolidine.

In the reaction of process d) any conventional peptide synthesis methods may be used.

The reaction may comprise the formation of, optionally *in situ*, an activated derivative of an acid, eg an anhydride or dicyclohexylcarbodiimide derivative. The reaction may be carried out in a solvent which is inert under the reaction conditions, eg dichloromethane or ethyl acetate, at a temperature of from -10°C to the boiling point of the solvent, preferably of from 0°C to 30°C. The reaction may be carried out in the presence of a base, eg triethylamine. When the reaction involves dicyclohexylcarbodiimide it may be carried out in the presence of an activating agent, eg hydroxybenzotriazole.

The reaction will of course vary with the particular activated derivative used.


In process e) the salts may be formed by reacting the free acid, or a salt, ester, amide or derivative thereof, or the free base, or a salt or derivative thereof, with one or more equivalents of the appropriate base or acid. The reaction may be carried out in a solvent or medium in which the salt is insoluble or in a solvent in which the salt is soluble, eg ethanol, tetrahydrofuran or diethyl ether, which may be removed *in vacuo*, or by freeze drying. The reaction may also be a metathetical process or it may be carried out on an ion exchange resin.

Pharmaceutically acceptable salts of the compounds of formula I include ammonium salts, alkali metal salts, eg sodium and potassium salts; alkaline earth metal salts, eg the calcium and magnesium salts; salts with organic bases, eg salts with dicyclohexylamine or N-methyl-D-glucamine; and salts with amino acids, eg with arginine, lysine etc. Also, when the molecule contains a basic group, salts with organic or inorganic acids, eg with HCl, HBr, H_2SO_4 , H_3PO_4 , methanesulfonic, toluenesulfonic, maleic, fumaric or camphorsulfonic acids. The non-toxic physiologically acceptable salts are preferred, although other salts are also useful, eg in isolating or purifying the product.

The starting materials for the above processes are either known or may be made from known compounds using conventional processes. Thus compounds of formula II in which n is 0 may be made by reaction of a compound of formula IV,




or a salt thereof,



or a salt, ester, amide or protected derivative thereof,

in which R_3 and Y are as defined above.

Compounds of formula II in which n is 1 may be made by reacting a compound of formula IV, or a salt thereof, with a compound of formula V,



or a salt, ester, amide or protected derivative thereof

in which X is as defined above,

for example in a solvent which is inert under the reaction conditions, eg benzene, at a temperature of from 0° to 100°C and preferably of from 0° to 25°C and in the presence of a base, eg 1,5-diazabicyclo [4.3.0] non-5-ene.

Compounds of formula III may be made from the appropriate acid or a derivative thereof using conventional processes known per se.

The compounds of formula I, and the intermediates therefor, may be isolated from their reaction mixtures using conventional techniques known per se.

The processes described above may produce the compound of formula I or a derivative thereof. It is also within the scope of this invention to treat any derivative so produced to liberate the free compound of formula I, or to convert one derivative into another.

in which R_3 and Y are as defined above,

with glyoxylic acid (or a salt, ester, amide or protected derivative thereof)

eg in an alkanol such as ethanol, at room temperature.

The compounds of formula II may exist in the tautomeric form of formula VII,

VII

In addition to the processes described above the compounds of formula I may be made by a variety of processes which are analogous to those known for the production of structurally similar compounds.

We further provide the compounds of formula II and salts, esters, amides and protected derivatives thereof, which are useful as intermediates.

Pharmaceutically acceptable esters include esters with C1 to 10 alcohols, eg alkyl C 1 to 6 esters and esters with benzyl alcohol. The amides may be, for example, unsubstituted or mono- or di-C 1 to 6 alkyl amides and may be made by conventional techniques, eg reaction of an ester of the corresponding acid with ammonia or an appropriate amine.


We prefer compounds of formula I in which D is a chain comprising from 2 to 11 atoms, more preferably from 3 to 11 and most preferably from 3 to 6 atoms.

We prefer the atoms in chain D to be selected from C and N. We further prefer that less than 4 N atoms are present in the chain, more preferably less than 3 and most preferably only one. When there is one N atom in the chain we prefer it to be less than 5 atoms away from the group $C=X$, more preferably less than 3 atoms away and most preferably one atom away.

The chain may optionally be substituted. We prefer such substituents to be selected from alkyl C 1 to 10, phenyl and aminoalkyl C 1 to 6. We prefer the substituents to be at each or either end of the chain D. Thus when the substituent is alkyl C 1 to 10 or aminoalkyl C 1 to 6 we prefer it to be at the end adjacent the group $C=X$. The alkyl substituent is preferably alkyl C 1 to 6, more prefer-

ably alkyl C 1 to 3 and most preferably methyl. The aminoalkyl substituent is preferably aminoalkyl C1 to 4 and more preferably aminobutyl. When the substituent is phenyl we prefer it to be at the end of the chain D remote from the group $C=X$.

We prefer the O or S containing substituent to be capable of chelating organic zinc. The O or S containing substituent is preferably at a position of from 3 to 5 atoms away from the group $C=X$, and more preferably 3 atoms away from the group



in which R_3 , Y and n are as defined above,

R is hydrogen, alkyl C1 to 10 or alkyl C1 to 6 substituted by NH_2 ,

Z is $R_2CH(COOH)NH$ - or R_2SCH_2 -,

R_1 is hydrogen or R_1CO -,

R_2 is alkyl C 1 to 10 or phenylalkyl C7 to 12, and

R_3 is alkyl C 1 to 10 or phenyl,

and pharmaceutically acceptable salts, esters and amides thereof.

We prefer those compounds of formula I in which Z is $R_2CH(COOH)NH$ -.

When Z is $R_2CH(COOH)NH$ - we prefer the partial structure $-NHCHRCON$ - in formula VIII to be part of a naturally occurring amino acid. We specifically provide compounds in which Z is $R_2CH(COOH)NH$ - and the two $-COOH$ groups are in different forms, eg where one is esterified and the other is not. We also prefer the group $COOH$ in the substituent Z to be in the form of an ester or amide, eg to be an alkyl C 1 to 6 ester, preferably an ethyl ester. We further prefer the carbon atom to which the group $COOH$ or its derivative in the substituent Z is attached to be in the S configuration.

Where any of R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , R_9 , R_{10} or R_{11} represent alkyl they may individually be straight, branched or cycloalkyl, eg containing up to and including 6 carbon atoms. We prefer R to be alkyl C 1 to 6 or aminoalkyl C 1 to 6. When R is unsubstituted alkyl C 1 to 6 we prefer R to be

$C=X$.. When the substituent contains O we prefer it to be a $C=O$ group and more preferably a $-COOH$ group or a derivative thereof, eg an alkyl C 1 to 6 ester. When the substituent contains S we prefer it to be a group $-SH$ or a protected derivative thereof, eg acetylthio.

X, is preferably O.

We particularly prefer the group of compounds of formula VIII,

VIII

methyl. When R is aminoalkyl C 1 to 6 we prefer the NH_2 group to be at the end of an unbranched chain, in particular we prefer R to be the group $-CH_2CH_2CH_2CH_2NH_2$. We further prefer the carbon atom to which R is attached to be in the S configuration.

When R_2 is alkyl we prefer it to be a straight chain alkyl, preferably a C 1 to 6 alkyl, most preferably n-propyl. When R_2 is phenylalkyl C7 to 12 we prefer the alkyl chain to comprise 1 to 3 carbon atoms. In particular, when R_2 is phenylalkyl, we prefer R_2 to be phenylethyl.

We prefer R_3 to be alkyl C 1 to 10, more preferably alkyl C1 to 6 and most preferably methyl.

We prefer R_{10} to be alkyl C 1 to 6 and more preferably methyl.

Where R_3 represents alkyl it may be straight, branched or cycloalkyl, eg containing up to and including 10 carbon atoms. The term cycloalkyl includes any mono-, bi- or tri-cyclic alkane. When R_3 represents a 5 membered heterocyclic group we prefer only one heteroatom to be present. When R_3 represents a 6 membered heterocyclic group we prefer one or two heteroatoms to be present, preferably selected from N and O. Examples of R_3 are methyl, pyridyl, furyl, methoxy, methylthio, t-butyl, isopropyl, cyclohexyl, morpholinyl, adamantyl, methylamino, benzyl, naphthyl and phenyl optionally substituted by methylthio, methoxy, methyl, ethyl, chloro or CF_3 .

We prefer R_4 to be alkyl C 1 to 10 or cycloalkyl C3 to 10, more preferably alkyl C 1 to 6 or cycloalkyl C3 to 6. We particularly prefer R_4 to be either t-butyl or cyclohexyl.

Y is preferably O, or more preferably S.

We prefer n to be 0.

We prefer the -COOH substituent on the Y containing heterocyclic ring to be underivatised. We further prefer the asymmetric carbon atom of the Y containing heterocyclic ring to be in the S configuration.

R₁ is preferably hydrogen.

We particularly prefer the specific group of compounds of formula VIII in which Z is R₂CH-(COOH)NH-, Y is S, R is methyl or aminobutyl, n is 0, R₂ is n-propyl or phenylethyl and R₃ is t-butyl and pharmaceutically acceptable salts, esters and amides thereof.

The preferred salts of the compounds of formula VIII are maleates, hydrochlorides, ammonium salts or dicyclohexyl-ammonium salts.

The compounds of formula I may contain one or more asymmetric carbon atoms and may therefore exhibit optical and/or diastereo isomerism. Diastereoisomers may be separated using conventional techniques, eg chromatography or fractional crystallisation. The various optical isomers may be isolated by separation of a racemic or other mixture of the compounds using conventional, e.g. fractional crystallisation or HPLC, techniques. Alternatively the desired optical isomers may be made by reaction of the appropriate optically active starting materials under conditions which will not cause racemisation. We prefer those compounds of formula I and formula VIII in which any asymmetric carbon atoms are in the S configuration.

The compounds of the invention are advantageous in that they are more efficacious, produce less side effects, are longer acting, more readily absorbed, less toxic, distributed in the body tissues in a different manner or have other advantageous properties when compared to compounds of similar structure.

The compounds of the invention are useful because they possess pharmacological properties. In particular they inhibit angiotensin converting enzyme and thus block conversion of the decapeptide angiotensin I to angiotensin II (see Example A). Angiotensin II is a potent vasoconstrictor in mammals. It also stimulates aldosterone release which results in salt and fluid retention. Increased blood pressure is the physiological result of these changes. Inhibitors of angiotensin converting enzyme are thus effective antihypertensive agents in a variety of animal models (see Example B) and are indicated for use clinically, for example, in patients with renovascular, malignant or essential hypertension or chronic congestive heart failure. See, for example, D W Cushman et al., *Biochemistry* 16, 5484 (1977) and E W Petrillo and M A Ondetti, *Med. Res. Rev.* 2 93 (1982).

Thus, the compounds of this invention are useful as antihypertensives in treating hypertensive mammals, including humans and they can be utilised to achieve reduction of blood pressure, eg in formulations containing appropriate pharmaceutically acceptable excipients, diluents or carriers. The compounds of the invention can be administered - (to animals or humans) in unit dosages of 1 to 500mg generally given several times, eg 1 to 4 times, per day thus giving a total daily dose of from 1 to 2000 mg per day. The dose will vary depending on the type and severity of disease, weight of patient and other factors which a person skilled in the art will recognise.

The compounds of this invention may be given in combination with other pharmaceutically active compounds, eg diuretics or antihypertensives. The dosage of the other pharmaceutically active compound can be that conventionally used when the compound is administered on its own, but is preferably somewhat lower. To illustrate these combinations, one of the antihypertensives of this invention effective clinically in the range, eg 1-200 milligrams per day, can be combined at levels ranging, eg from 1-200 milligrams per day with the following antihypertensives and diuretics in dose ranges per day as indicated:

hydrochlorothiazide (15-200mg), chlorothiazide (125-2000mg), ethacrynic acid (15-200mg), amiloride (5-20mg), furosemide (5-80mg), propranolol (20-480mg), timolol (5-50mg) nifedipine - (20-100mg), verapamil (120-480mg) and methyldopa (65-2000mg). In addition, the triple drug combinations of hydrochlorothiazide (15-200mg) plus amiloride (5-20mg) plus converting enzyme inhibitor of this invention (1-200mg) or hydrochlorothiazide (15-200mg) plus timolol (5-50mg), plus the converting enzyme inhibitor of this invention (1-200mg) are contemplated. The above dose ranges may be adjusted on a unit basis as necessary to permit divided daily dosage. Also, the dose may vary depending on the severity of the disease, weight of patient and other factors which a person skilled in the art will recognise.

According to our invention we also provide a pharmaceutical composition comprising preferably less than 80%, more preferably less than 50%, eg 1 to 20%, by weight of a compound of formula I, or a pharmaceutically acceptable salt or ester thereof, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier.

Thus the compound may be put up as a tablet, capsule, dragee, suppository, suspension, solution, injection, implant, a topical, eg transdermal, preparation such as a gel, cream, ointment, aerosol or a polymer system, or an inhalation form, eg an aerosol or a powder formulation.

We prefer compositions which are designed to be taken oesophageally and to release their contents in the gastrointestinal tract. Thus we prefer tablets which may, for example, be made by direct compression. In such a process the active ingredient is mixed with one or more of modified forms of starch, calcium phosphate, a sugar eg lactose, microcrystalline cellulose and/or other directly compressible excipients, together with lubricant(s), eg stearic acid or magnesium stearate, flow aid(s), eg talc or colloidal silicon dioxide, and disintegrant(s), eg starch or the materials sold under the Trade Marks, Nymcel, Ac-Di-Sol, Explotab and Plasdone XL. Tablets are then formed by direct compression, and may be sugar or film coated eg with hydroxypropylmethylcellulose.

Alternatively the active ingredient may be granulated before tableting. In such cases the active ingredient is mixed with one or more of starch, calcium phosphate, a sugar eg lactose, microcrystalline cellulose or other suitable excipients and granulated with a binder such as starch, pregelled starch, polyvinylpyrrolidone, gelatine, a modified gelatine, or a cellulose derivative, eg hydroxypropylmethylcellulose. The mass is then dried, sieved and mixed with lubricant(s), flow aid(s) and disintegrant(s), such as described in the previous paragraph. Tablets are then formed by compression of the granules, and may be sugar or film coated, eg with hydroxypropylmethylcellulose.

As a further alternative a powder, blend or granules, such as those described above as intermediates in tableting, may be filled into a suitable, eg gelatine, capsule.

In order to improve the bioavailability, or decrease variability of availability, of the active ingredient the compound may be:-

a) dissolved in a suitable solvent, eg polyethylene glycol, Gelucaire, arachis oil, a - (hydrogenated) vegetable oil or beeswax and the solution is then filled into a gelatine capsule,

b) produced as a spray-dried or freeze-dried form prior to mixing with other excipients,

c) milled and/or micronised to produce a powder with a large surface area prior to mixing with other excipients,

d) made into a solution and distributed over an inert excipient having a large surface area, eg colloidal silicon dioxide. The solvent is evaporated and further excipients added,

e) formed into a complex with cyclodextrin prior to mixing with other excipients. This complex also assists in increasing light stability, or

f) made into a solid solution or co-precipitated, eg with polyvinylpyrrolidone, polyethyleneglycol, modified cellulose, hydroxypropylmethylcellulose, urea or a sugar prior to mixing with further excipients.

The compounds, either in their normal form or in a modified form, eg as described immediately above, may be formulated in a controlled release form. Thus the compound may be dispersed, or contained in, a polymer matrix formed from, for example, ethylcellulose, hydroxypropylmethylcellulose or the product sold under the Trade Mark Eudragit. Alternatively the compound may be formulated as a tablet or beads which are surrounded by a semi-permeable membrane, eg shellac, ethylcellulose or an acrylate/methacrylate polymer.

Certain of the compounds of formula I can form hydrates or solvates, eg with an alcohol such as ethanol or, for example when Y is NH can exist in tautomeric forms.

The invention is illustrated, but in no way limited by the following Examples in which temperatures are in degrees centigrade.

Example 1

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-(S)-carboxylic acid

a) Benzyl 2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-carboxylate

A solution of benzenecarbothioic acid hydrazide (2g) and benzyl glyoxylate (2.6g) in ethanol (5ml) was stirred at room temperature for 18 hours under nitrogen. The solvent was removed by evaporation and the residue was flash chromatographed to yield the sub-title product - (3.5g) as a beige solid.

A mass spectrum showed M^+ 298 (base peak 163).

$C_{18}H_{18}N_2O_2S$ requires MWt 298.

b) Benzyl 3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-(R)-carboxylate

A stirred mixture of N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanine (3.1g) and 1-hydroxybenzotriazole (1.7g) in dichloromethane (100ml) was treated with a solution of the product of step a) - (6.85g) in dichloromethane (25ml). A solution of dicyclohexylcarbodiimide (2.26g) in dichloromethane (20ml) was added over 20 minutes and the mixture was stirred at room temperature for 2 days under nitrogen.

The suspended solid was filtered, the filtrate evaporated and the residue was purified by flash chromatography to give the sub-title product - (5.23g) as a gum.

A fast atom bombardment mass spectrum showed M^+560 (base peak 91).

$C_{21}H_{25}N_3O_3S$ requires MWt 559.

c) Benzyl 3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-(S)-carboxylate

A solution of the product from step b) (0.16g), pyrrolidine (0.16ml) and 3A molecular sieves - (0.2g) in acetonitrile (3.2ml) was stirred at room temperature for 3.5 hours. The mixture was poured into water and extracted with ether, dried over magnesium sulphate and evaporated. The residue was flash chromatographed to give the sub-title product (0.05g) as a gum.

A fast atom bombardment mass spectrum showed M^+560 (base peak 91).

$C_{21}H_{25}N_3O_3S$ requires MWt 559.

d) 3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-(S)-carboxylic acid

A solution of the product from step c) (0.26g) in ethanol (20ml) was treated with 10% palladium on charcoal (0.1g) and stirred in a pressure vessel under hydrogen at 3 atmospheres at room temperature for 3 days. The catalyst was filtered off and the filtrate evaporated. The residue was triturated with ether to give the title product (0.08g) as a white solid, m.p. $180.5^\circ-182^\circ$.

A mass spectrum (FAB) showed M^+470 (base peak 234).

$C_{20}H_{23}N_3O_3S$ requires MWt 469.

Example 2

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-(R)-carboxylic acid

A solution of the product from Example 1, step b) (0.43g) in ethanol (100ml) was treated with 10% palladium on charcoal (0.1g) and stirred in a pressure vessel under hydrogen at 3 atmospheres at room temperature for 3 days. The catalyst was filtered off and the filtrate evaporated. The residue was triturated with a mixture of ether and petroleum ether (bp $40^\circ-60^\circ$) to give the title product (0.19g) as a pale grey, non-crystalline solid.

A mass spectrum (FAB) showed M^+470 (base peak 234).

$C_{20}H_{23}N_3O_3S$ requires MWt 469.

Example 3

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-phenyl-1,3,4-thiadiazole-2-carboxylic acid

a) Ethyl 2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-carboxylate

A solution of benzenecarbothioic acid hydrazide (0.4g) and ethyl glyoxylate (0.4g) in ethanol (1ml) was stirred at room temperature for 2 hours. The solvent was removed by evaporation and the residue re-evaporated with toluene (x2) to yield the sub-title product (0.7g) as a gum.

A mass spectrum showed M^+236 (base peak 163).

$C_{11}H_{13}N_3O_2S$ requires MWt 236.

b) Ethyl 3-(3-acetylthio-1-oxopropyl)-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-carboxylate

A solution of the product of step a) (2.36g) in toluene (100ml) was treated with polyvinylpyridine - (2.0g) and 3-acetylthiopropionyl chloride (1.7g) and the mixture stirred at room temperature for 4 hours. The mixture was filtered and the filtrate stirred with a saturated solution of sodium bicarbonate (100ml) for 1 hour. The organic phase was separated, washed with water, dried and evaporated to a gum. The residue was purified by flash chromatography to give the sub-title product (2.62g) as an oil.

A mass spectrum showed M^+366 (base peak 163).

$C_{20}H_{21}N_3O_3S_2$ requires MWt 366.

c) 2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-phenyl-1,3,4-thiadiazole-2-carboxylic acid

A solution of the product of step b) (2.6g) in methanol (20ml) was cooled to 0° under nitrogen and treated dropwise with a solution of potassium hydroxide (1.42g) in water (8ml). The mixture was allowed to warm to room temperature over 2 hours and then partitioned between ethyl acetate and water. The aqueous phase was acidified with 2N HCl and the organic phase separated, washed with water and dried. Evaporation yielded an oil which slowly crystallised to give the title product (0.7g) as white crystals. mp $145-6^\circ$.

Found: C 48.54, H 4.17, N 9.49, S21.68%

$C_{12}H_{12}N_2O_2S_2$ requires

C 48.65, H 4.05, N 9.46, S21.62%

Example 4

5-t-Butyl-3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid

a) Benzyl 5-t-butyl-2,3-dihydro-1,3,4-thiadiazole-2-carboxylate

A solution of t-butylcarbothioic acid hydrazide - (0.7g) and benzyl glyoxylate (1g) in ethanol (15ml) was stirred under nitrogen for 16 hours. The solvent was removed by evaporation and the residue was purified by flash chromatography (petroleum ether/ethyl acetate eluent) to yield the sub-title product (1.1g) as a gum.

b) Benzyl 5-t-butyl-3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(R)-carboxylate

A stirred mixture of N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanine (0.57g) and 1-hydroxybenzotriazole (0.28g) in dichloromethane (40ml) was treated with a solution of the product of step a) - (1.14g) in dichloromethane (5ml). Dicyclohexylcarbodiimide (0.42g) was added and the mixture stirred at room temperature for 16 hours under nitrogen. The suspended solid was removed by filtration and the filtrate evaporated to a gum. The residue was purified by flash chromatography to give the sub-title product (0.82g) as an oil.

A mass spectrum (FAB) showed M^+ 540 (base peak 91).

$C_{23}H_{27}N_3O_5S$ requires MWt 539.

c) Benzyl 5-t-butyl-3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylate

A solution of the product from step b) (1.0g) and pyrrolidins (1g) in dry acetonitrile (30ml) was treated with crushed 3A molecular sieves and the mixture stirred at room temperature for 6 hours. The volatile materials were removed by evaporation and the S,S,S isomer separated from the more polar S,S,R isomer by flash chromatography - (petroleum ether/ethyl acetate eluent). The sub-title product (0.4g) was isolated as a clear gum

A mass spectrum showed M^+ 539 (base peak 234).

$C_{23}H_{27}N_3O_5S$ requires MWt 539.

d) 5-t-Butyl-3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid

A solution of the product from step c) (0.67g) in ethanol (100ml) was treated with 10% palladium on carbon (0.6g) and the mixture stirred under 1 atmosphere of hydrogen for 16 hours. The catalyst was removed by filtration and the filtrate reduced in volume to ca. 2ml by evaporation. The cooled solution yielded the title product (0.3g) as white crystals, mp 165-8°.

Found: C,58.87; H,6.89; N,9.34; S,7.21%

$C_{22}H_{21}N_3O_5S$ requires

C,58.80; H,6.90; N,9.35; S,7.13%

A mass spectrum (FAB) showed M^+ 450 (base peak 234).

$C_{22}H_{21}N_3O_5S$ requires MWt 449.

Example 5

5-t-Butyl-3-[N²-(1-(S)-carboxy-3-phenylpropyl)-L-lysyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid

a) Benzyl 2-hydroxy-4-phenylbutanoate

A solution of 2-hydroxy-4-phenylbutanoic acid (20.4g), triethylamine (15.9ml) and benzyl bromide (12.75ml) in ethyl acetate (64ml) was heated under reflux for 16 hours. The solution was cooled and poured into a mixture of water and ether. The separated organic extract was washed with saturated sodium bicarbonate solution and water, dried over magnesium sulphate and filtered. The filtrate was evaporated and the residue purified by flash chromatography (petroleum ether/ethyl acetate eluent) to yield the sub-title product as a yellow oil (14g).

A mass spectrum showed M^+ 270 (base peak 91).

$C_{17}H_{19}O_3$ requires MWt 270.

b) N⁶-Benzyloxycarbonyl-N²-(1-(S)-benzyloxycarbonyl-3-phenylpropyl)-L-lysine hydrochloride

A solution of the product from step a) (13.8g) and pyridine (6.6ml) in dichloromethane (136ml) was added over 0.5 hours under nitrogen to a stirred solution of trifluoromethanesulphonic anhydride (12.9ml) in dichloromethane (136ml) cooled to 5°C. After a further 0.5 hours the solution was washed with water, dried over magnesium sulphate, filtered and the filtrate evaporated.

The residue was taken up in dichloromethane - (136ml) and added to a solution of N⁶-benzyloxycarbonyl-L-lysine t-butyl ester (15.5g) and triethylamine (6.5ml) in dichloromethane - (136ml). The mixture was stirred at room temperature for 1 hour, heated under reflux for 2.5 hours, cooled, washed with water, dried over magnesium sulphate and filtered. The filtrate was evaporated and the residue purified by flash chromatography - (ether/petroleum ether eluent) to separate and isolate the more polar SS isomer.

A solution of the SS t-butyl ester (0.5g) in ether (15ml) was cooled to +5° and saturated with hydrogen chloride for 2 hours. The solution was stirred at room temperature for a further 18 hours and the solvent was then removed by evaporation. Trituration of the residue in ether gave the sub-title product as a white solid (0.39g).

A fast atom bombardment mass spectrum showed M⁺533 (base peak 91).

C₂₁H₂₄N₂O₆ requires MWt 532.

c) Benzyl 3-[N⁶-benzyloxycarbonyl-N²-(1-(S)-benzyloxycarbonyl-3-phenylpropyl)-L-lysyl]-5-t-butyl-2,3-dihydro-1,3,4-thiadiazole-2-(R)-carboxylate

A stirred solution of the SS product from step b) (5.88g) and 1-hydroxybenzotriazole (1.35g) in dichloromethane (85ml) was treated with a solution of the product of Example 4, step a) (5.87g) in dichloromethane (60ml). A solution of dicyclohexylcarbodiimide (2.1g) of dichloromethane (85ml) was added over 5 minutes and the mixture was stirred at room temperature for 18 hours under nitrogen. Triethylamine (1.4ml) was added and the suspended solid removed by filtration. The filtrate was evaporated and the residue purified by flash chromatography to give the sub-title product as an oil (2.1g).

A fast atom bombardment mass spectrum showed M⁺793 (base peak 91).

C₄₆H₄₂N₄O₈S requires MWt 792.

d) Benzyl 3-[N⁶-benzyloxycarbonyl-N²-(1-(S)-benzyloxycarbonyl-3-phenylpropyl)-L-lysyl]-5-t-butyl-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylate

A solution of the product of step c) (2.1g) and pyrrolidine (1.6ml) in dry acetonitrile (60ml) was treated with crushed 3A molecular sieves and the mixture stirred at room temperature for 24 hours under nitrogen. The volatile materials were removed by evaporation and the SSS isomer separated from the more polar SSR isomer by flash chromatography. The SSS sub-title product - (0.47g) was isolated as a clear oil.

A fast atom bombardment mass spectrum showed M⁺793 (base peak 91).

C₄₆H₄₂N₄O₈S requires MWt 792.

e) 5-t-Butyl-3-[N²-(1-(S)-carboxy-3-phenylpropyl)-L-lysyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid

A solution of the product from step d) (1.1g) in ethanol (90ml) was treated with 10% palladium on carbon (0.9g) and the mixture stirred under 1 atmosphere of hydrogen for 1 hour. The catalyst was removed by filtration and the filtrate evaporated. The residue was recrystallised from a mixture of tetrahydrofuran and ethanol to give the title product as a white solid (0.24g)

mp slowly decomposes at 180 -190°

Found: C 55.86 H 6.97 N 11.24 S 6.56 H₂O 2.83

C₂₂H₂₄N₄O₆S. 0.77H₂O

Requires: C 56.11 H 7.23 N 11.39 S 6.51 H₂O 2.82

A fast atom bombardment mass spectrum showed M⁺479 (base peak 84).

C₂₂H₂₄N₄O₆S requires MWt 478.

Example 6

5-t-Butyl-3-[N-(1-(S)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid

a) Ethyl 2-([(trifluoromethyl)sulphonyl]oxy)-pentanoate

Under nitrogen, a solution of pyridine (11.9g) in dry dichloromethane (500ml) was rapidly stirred at -22° while trifluoromethane sulphonic anhydride - (40.5g) was added dropwise. After the addition, the white slurry was stirred at -22° for 15 minutes and then a solution of ethyl 2-hydroxy pentanoate - (16.8g) in dichloromethane was added over 2 minutes at this temperature. The temperature was then allowed to rise to room temperature and the mixture was stirred vigorously for 1 hour, after which time the white solid was filtered off, washed well with dichloromethane and the combined washings

and filtrate were evaporated to dryness. The oily solid obtained was taken up in 60 -80° petroleum ether and was passed down a short pad of silica, eluting with more petroleum ether. The petroleum ether solution was evaporated to dryness leaving an oil (23.4g)

NMR, CDCl_3 , δ : 1.0(3H,t), 1.32(3H,t), 1.5(2H,m), 2.0(2H,m), 4.3(2H,m) 5.12(1H,t)

b) N-(1-Ethoxycarbonylbutyl)-L-alanine benzyl ester

L-Alanine benzyl ester hydrochloride (10.0g) was converted to the free-base in dichloromethane using triethylamine. The resulting mixture was evaporated to dryness and the residue was slurried with several portions of diethyl ether. The combined ether solutions were evaporated to dryness leaving the free-base as an oil.

R,S

NMR CDCl_3 , δ , 0.9 (3H,t), 1.24 (3H,t), 1.30 (3H,d), 1.38 (2H,m), 1.6 (2H,m), 3.27 (1H,t), 3.4 (1H,q), 4.12 (2H,m), 5.15 (2H,q), 7.35 (5H,s).

S,S

NMR CDCl_3 , δ : 0.9 (3H,t); 1.28 (3H,t); 1.38 (3H,d); 1.3-1.8 (4H,m); 3.28 (1H,t); 3.4 (1H,q); 4.18 (2H,m); 5.17 (2H,q); 7.35 (5H,s).

Found: C, 55.62; H, 8.47; N, 6.32%

c) N-(1-(S)-Ethoxycarbonylbutyl)-L-alanine

The S,S diastereoisomer benzylester from step b) (6.2g) in ethanol (250ml) was hydrogenated at 3 atmospheres at room temperature for 30 minutes over 10% palladium on charcoal (0.6g). The catalyst was removed by filtration and the filtrate was evaporated to near dryness. The residue was slurried with diethyl ether and the white solid was filtered off and was dried to give the required product (3.8g) mp 153 -4°.

The oil was dissolved in dichloromethane - (200ml) and triethylamine (7.2g) was added. The resulting solution was stirred under nitrogen at room temperature while a solution of the product from step a) (12.8g) in dichloromethane was added dropwise over 30 minutes. The resulting mixture was stirred for 2 hours at room temperature, for 2 hours at reflux and then was evaporated to dryness leaving an oil residue which was purified by flash chromatography on silica eluting with 60 -80° petroleum ether/diethyl ether 5:1 to give the R,S, - (4.8g, 34%) and S,S, (4.7g, 33%) diastereoisomer in order of elution

$\text{C}_{10}\text{H}_{17}\text{NO}_4$ requires

C, 55.29; H, 8.75; N, 6.45%

d) Benzyl 5-t-butyl-3-[N-(1-(S)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(R)-carboxylate

Under nitrogen at room temperature a mixture of the S,S amino acid from step c) (0.62g) and 1-hydroxybenzotriazole (0.45g) in dry dichloromethane (125ml) was stirred for 30 minutes

with benzyl 5-t-butyl-2,3-dihydro-1,3,4-thiadiazole-2-carboxylate (1.6g). Dicyclohexylcarbodiimide - (0.6g) was then added and the resulting mixture was stirred for 18 hours, filtered and the filtrate evaporated to dryness. The residue was purified by column chromatography on silica eluting with diethyl ether/petroleum ether (60 -80°), 1:1 to give the required diester as an oil (1.1g)

NMR CDCl₃,delta: 0.9(3H,t), 1.2-1.7(19H,m), 3.3-(1H,t) 4.2(3H,m), 5.2(2H,q), 6.18(1H,s) 7.35(5H,s).

e) Benzyl 5-t-butyl-3-[N-(1-(S)-ethoxycarbonyl-butyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylate

Under nitrogen, pyrrolidine (1.5ml) was added to a solution of the 'S,S,R' ester (step d) (1.6g) and the resulting solution was stirred at room temperature for 24 hours. The 1:1 mixture of S,S,R' and S,S,S esters so produced was separated by flash chromatography on silica eluting with ethyl acetate/petroleum ether 60 - 80°, 1:3 to give 0.65g of each isomer. The S,S,R isomer was recycled such that the total conversion was 81%. NMR CDCl₃,delta: 0.9(3H,t), 1.2-1.7(19H,m), 3.3(1H,t) 4.2-(3H,m), 5.17(2H,s), 6.18(1H,s)

7.35(5H,s).

f) 5-t-Butyl-3-[N-(1-(S)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid

The S,S,S benzyl ester from step e) (1.8g) in ethanol (500ml) was hydrogenated over 10% Pd on charcoal (1.8g) at atmospheric pressure and room temperature for 5 hours. The catalyst was removed by filtration and the filtrate was evaporated to dryness. The residue was triturated with a 1:1 mixture of ether/petroleum ether 60 -80° to give the required acid as a white solid (1.3g). mp 183 -5°.

Found: C, 47.13; H, 7.89; N, 9.31; S, 7.12%

C₁₇H₂₈N₂O₅S. 2.5 H₂O requires

C, 47.22; H, 7.87; N,9.72; S, 7.41%

Example 7

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-[4-(methylthio)phenyl]-1,3,4-thiadiazole-2-(S)-carboxylic acid

5 a) 1-[(4-(Methylthio)phenyl)thioxomethyl]pyrrolidine

A mixture of 4-(methylthio)benzaldehyde - (50.0g) and sulphur (15.8g) was cooled to 0° and pyrrolidone (41.1ml) was added over 30 minutes. On complete addition the whole was heated under reflux for 1.5 hours. The mixture was poured, whilst warm, into ethanol (250ml) and the resulting solid filtered off. Recrystallization from ethanol afforded the sub-title compound as a fawn, crystalline solid, (71.3g). mp 116.5-118°.

b) 4-[4-(Methylthio)phenyl]-4-(pyrrolidinium-1-ylidene)-3-thiobutanoic acid bromide

A solution of the product of step a) (20.0g) and bromoacetic acid (12.9g) in benzene (100ml) was stirred at room temperature under nitrogen for 18 hours. The resulting precipitate was filtered off and washed with ether to yield the sub-title compound - (28.6g) as a white solid. mp 157-158°.

30 c) [4-(Methylthio)phenyl]thioxomethylthio]acetic acid

Hydrogen sulphide was passed through a solution of the product of step b) (25.0g) in methanol - (250ml) and cooled in an ice bath, for a period of 3 hours.

After standing at 0° for 18 hours the solvent was removed under reduced pressure and the residue was triturated with water. The solid was filtered off and recrystallised from petroleum ether to afford the sub-title compound (16.7g) as a red crystalline solid. mp 117°.

45 d) 4-(Methylthio)phenylcarbothioic acid hydrazide

To a solution of the product of step c) (15.0g) in methanol (200ml) was added aqueous potassium hydroxide (58.0ml, 1M) followed by hydrazine monohydrate (3.1ml) dropwise over 30 minutes. After stirring at room temperature for 1 hour the mixture was acidified to pH 5 with concentrated hydrochloric acid. The resulting precipitate was filtered off and recrystallised from ethanol to afford the sub-title compound (9.9g) as pale yellow plates. mp 152-153°.

e) t-Butyl 2,3-dihydro-5-[4-(methylthio)phenyl]-1,3,4-thiadiazole-2-carboxylate

The product of step d) (3.0g) was stirred at room temperature under nitrogen with t-butyl glyoxylate (2.0g) in methanol (100ml) for 18 hours. The mixture was evaporated and the residue purified by flash chromatography (ether/petroleum ether 1:3) to afford the sub-title compound (3.9g) as a pale yellow solid. mp 71-72°.

f) t-Butyl 3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-[4-(methylthio)phenyl]-1,3,4-thiadiazole-2-(S)-carboxylate

Prepared from the product of step e) and N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanine by similar processes to those of Example 1, steps b and c as a yellow oil.

A mass spectrum (FAB) showed M⁺572 (base peak 234)

C₂₈H₃₂N₂O₅S₂ requires 571.

g) 3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-[4-(methylthio)phenyl]-1,3,4-thiadiazole-2-(S)-carboxylic acid

A solution of the product (1.0g) from step f) in dry diethyl ether (50ml) was saturated with hydrogen chloride for 3 hours. The solvent was evaporated and the residue was purified by chromatography to give the title compound (0.1g) as a pale yellow solid. mp 163-164°

Found: C, 57.46; H, 5.53; N, 8.01; S, 12.14

C₂₈H₃₂N₂O₅S₂. 0.5 H₂O requires

C, 57.25; H, 5.72; N, 8.01; S, 12.21

Example 8

Ammonium 2,3-dihydro-3-(3-mercapto-2-(S)-methyl-1-oxopropyl)-5-phenyl-1,3,4-thiadiazole-2-(S)-carboxylate monohydrate

a) Benzyl 3-(3-acetylthio-2-(S)-methyl-1-oxopropyl)-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-carboxylate

3-Acetylthio-2-(S)-methylpropanoyl chloride - (2.3g) in dichloromethane (30ml) was added over 5 minutes to a stirred mixture of the product of Example 1, step a) (3.6g) and polyvinylpyridine - (2.4g) in dichloromethane (60ml). The mixture was stirred at room temperature for 20 hours and 3-

acetylthio-2-(S)-methylpropanoyl chloride (1.2g) was then added. The mixture was stirred for a further 2 hours, filtered and the filtrate stirred with a saturated solution of sodium bicarbonate for 1 hour. The organic phase was separated, washed with water, dried and evaporated to a gum. The residue was purified by flash chromatography to give the sub-title product (4.4g) as an oil.

b) Ammonium 2,3-dihydro-3-(3-mercapto-2-(S)-methyl-1-oxopropyl)-5-phenyl-1,3,4-thiadiazole-2-(S)-carboxylate

A solution of the product of step a) (2.8g) in methanol (50ml) under nitrogen was treated dropwise with a solution of potassium hydroxide (1.3g) in water (100ml). The mixture was stirred for a further 4 hours and then partitioned between ether and water. The separated aqueous phase was acidified with 2N HCl and extracted with ether. The organic phase was washed with water, dried and evaporated to an oil. The residue was purified by reverse phase HPLC to give the title product - (0.023g) as a white solid. mp 194-7°.

A mass spectrum showed M⁺310 (base peak 163).

C₂₇H₂₈N₂O₅S₂ Requires MWt 310.

Example 9

2-Cyclohexyl-5,6-dihydro-4-(3-mercapto-1-oxopropyl)-4H-1,3,4-thiadiazine-5-carboxylic acid

a) Benzyl 2-cyclohexyl-5,6-dihydro-4H-1,3,4-thiadiazine-5-carboxylate

A solution of 2-bromoprop-2-enoic acid benzyl ester (1.68g) in dry benzene (10.2ml) was added to cyclohexane carbothioic acid hydrazide (1.0g) in dry dimethylformamide (10ml). The mixture was cooled to 0° and stirred, under an atmosphere of nitrogen, during the gradual addition of 1,5-diazabicyclo[4.3.0]non-5-ene (0.78g) and then for a further 20 minutes at 0°. Ethyl acetate (100ml) was added and the mixture was washed with brine and dried over magnesium sulphate. The solvent was evaporated under reduced pressure and the resulting oil purified by flash chromatography using 10% ethyl acetate/90% petroleum ether as eluent to yield the sub-title product (1.0g) as a pink solid.

A mass spectrum showed M⁺318 (base peak 91).

C₂₇H₂₈N₂O₅S requires MWt 318.

b) Benzyl 4-(3-acetylthio-1-oxopropyl)-2-cyclohexyl-5,6-dihydro-4H-1,3,4-thiadiazine-5-carboxylate

3-Acetylthiopropionyl chloride (0.45g) and poly-(4-vinylpyridine) (0.8g) were added to a solution of the product of step a) (0.86g) in dry toluene (20ml). The mixture was stirred under an atmosphere of nitrogen for 20 hours. Diethyl ether (30ml) was added and the mixture filtered. The filtrate was evaporated and the product purified by flash chromatography to yield the sub-title product (1.05g) as a pale yellow oil.

Mass spectrum (FAB) showed M⁺449 (base peak 91).

C₂₂H₂₈N₂O₃S₂ requires MWt 448.

c) 2-Cyclohexyl-5,6-dihydro-4-(3-mercapto-1-oxopropyl)-4H-1,3,4-thiadiazine-5-carboxylic acid

1M Potassium hydroxide solution in methanol (5.83ml) was added to a solution of the product of step b) (0.87g) in methanol (10ml) and water (5ml). The mixture was stirred under an atmosphere of nitrogen for 2 hours. Acetic acid was added and the solvent evaporated under reduced pressure. The mixture was purified by flash chromatography using 1% acetic acid/ethyl acetate as eluent to yield the title compound (0.24g) as a white solid. mp 95-97°.

Mass spectrum showed M⁺316 (base peak 156).

C₁₈H₂₆N₂O₃S₂ requires MWt 316.

Example 10

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-phenyl-1,3,4-oxadiazole-2-carboxylic acid

a) Ethyl (benzoylhydrazono)acetate

A solution of benzoyl hydrazine (1.4g) and ethyl glyoxylate (1.32g) in ethanol (50ml) was stirred at room temperature for 24 hours. The solvent was evaporated and the residue treated with ether to give the sub-title product (1.8g) as white solid. mp 140-3°.

b) 3-(Acetylthio)propionic anhydride

A solution of 3-(acetylthio)propionic acid (3.4g) in ether (20ml) was treated dropwise with a solution of dicyclohexylcarbodiimide (2.1g) in ether with water-bath cooling. The mixture was stirred for 1.5 hours, filtered and the filtrate evaporated to give the sub-title product (3.2g) as a yellow oil.

c) Ethyl 3-(3-acetylthio-1-oxopropyl)-2,3-dihydro-5-phenyl-1,3,4-oxadiazole-2-carboxylate

A mixture of the product from step a) (2.6g) and the crude product from step b) (3.2g) in pyridine (0.9ml) was heated at 100° for 18 hours. The mixture was poured into water and extracted with ethyl acetate. The separated organic extract was washed with water, saturated aqueous sodium bicarbonate solution, water, dried and evaporated.

The residue was purified by flash chromatography to give the sub-title product (1.6g) as a yellow oil.

A mass spectrum showed M⁺350 (base peak 147).

C₁₈H₁₈N₂O₃S requires MWt 350.

d) 2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-phenyl-1,3,4-oxadiazole-2-carboxylic acid

A solution of the product from step c) (1.44g) in methanol (40ml) was cooled to 15° under nitrogen and treated dropwise with a solution of potassium hydroxide (0.69g) in water (40ml). The mixture was stirred at room temperature for 2 hours and then the solvents were evaporated. The residue was taken up in water and washed with ether. The aqueous phase was acidified with 2N HCl and extracted with ethyl acetate. The separated organic phase was washed with water, dried and evaporated. The residue was purified by flash chromatography to give a pale yellow solid. The solid was taken up in dichloromethane, treated with charcoal, filtered and the filtrate evaporated to give the title product (0.24g) as an off-white solid. mp 106-9°.

Example 11

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-[4-(trifluoromethyl)phenyl]-1,3,4-thiadiazole-2-carboxylic acid

a) 4-(Trifluoromethyl)phenylcarbothioic acid hydrazide

[[[4-(Trifluoromethyl)phenyl]thioxomethyl]thio]-acetic acid (6.7g) was dissolved in methanol - (50ml). Potassium hydroxide (1.34g) in water (15ml) was added followed by hydrazine hydrate - (1.28ml). The mixture was stirred for 4 hours at room temperature. Glacial acetic acid was added until the pH was 5 and the solvent was removed under reduced pressure. The product was extracted with diethyl ether (250ml) and the solvent was evaporated under reduced pressure. The resulting solid was crystallised from cyclohexane as pale pink crystals (4.0g) mp 114-115.5°.

b) Benzyl 2,3-dihydro-5-[4-(trifluoromethyl)phenyl]-1,3,4-thiadiazole-2-carboxylate

Benzyl glyoxalate (0.8g) was added to a solution of the product from step a) (1.0g) in dry ethanol (30ml). The mixture was stirred at room temperature under an atmosphere of nitrogen for 5 hours. The solvent was removed under reduced pressure and the product was crystallised from ethanol to yield the sub-title product (1.4g) as white crystals. mp 99-100.5°.

c) Benzyl 3-[3-acetylthio-1-oxopropyl]-2,3-dihydro-5-[4-(trifluoromethyl)phenyl]-1,3,4-thiadiazole-2-carboxylate

3-Acetylthiopropionyl chloride (0.36g), poly (4-vinyl pyridine) (0.8g) and the product from step b) - (0.8g) were stirred together in dry toluene (30ml) under an atmosphere of nitrogen for 20 hours. Diethyl ether was added and the solid was filtered off and washed with diethyl ether. The filtrate was evaporated under reduced pressure and the resulting product was crystallised from ethanol to yield the sub-title product (0.9g) as a white solid. mp 120-121°.

d) 2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-[4-(trifluoromethyl)phenyl]-1,3,4-thiadiazole-2-carboxylic acid

Potassium hydroxide in methanol (1M, 4.9ml) was added to a solution of the product from step c) (0.81g) in methanol (10ml) and water (5ml). The mixture was stirred under an atmosphere of nitro-

gen for 2 hours. Glacial acetic acid was added and the solvent was removed under reduced pressure. The product was purified by flash chromatography using 1% acetic acid/99% ethyl acetate as eluent to yield the title compound (0.23g) as a fawn solid. mp softens 93-75°.

A mass spectrum showed M⁺ 364 (base peak 231)

C₁₅H₁₁F₃N₂O₃S₂ requires MWt 364.

Example 12

Benzyl 4-[3-acetylthio-1-oxopropyl]-5,6-dihydro-1-methyl-2-phenyl-4H-1,3,4-triazine-5-carboxylate

a) Benzyl 5,6-dihydro-1-methyl-2-phenyl-4H-1,3,4-triazine-5-carboxylate

Benzenecarboximidic acid-N-methyl hydrazide monohydroiodide (0.43g) in dry dimethylformamide (10ml) and benzyl 2-bromo-prop-2-enoate (0.41g) in benzene (2.5ml) were cooled to 0° and stirred together under an atmosphere of nitrogen. A solution of 1,5-diazabicyclo[4.3.0]non-5-ene (0.4ml) in dimethylformamide (5ml) was added gradually and the mixture was stirred at 0° for a further 20 minutes. Ethyl acetate (100ml) was added and the mixture was washed with brine (30ml) and dried over magnesium sulphate. The solvent was removed at reduced pressure to yield the sub-title product (0.46g) as an orange oil.

A mass spectrum showed M⁺ 309 (base peak 91).

A fast atom bombardment mass spectrum showed M⁺ 310 (base peak 91).

C₁₅H₁₃N₃O₂ requires MWt 309.

b) Benzyl 4-[3-acetylthio-1-oxopropyl]-5,6-dihydro-1-methyl-2-phenyl-4H-1,3,4-triazine-5-carboxylate

3-Acetylthiopropionyl chloride (0.25g), poly(4-vinyl pyridine) (0.4g) and the product from step a) were stirred together in dry toluene (20ml) under an atmosphere of nitrogen for 24 hours. Diethyl ether (20ml) was added and the solid was filtered off and washed with diethyl ether. The filtrate was evaporated under reduced pressure. The resulting oil was purified by flash chromatography using 10% ethyl acetate/90% petroleum ether as eluent to yield the title compound (0.11g) as an amber oil.

A mass spectrum showed M⁺ 439 (base peak 93).

C₂₃H₂₃N₃O₃S requires MWt 439.

NMR CDCl₃ delta: 1.84(S,3H), 2.31(S), 2.35-3.35-(m), 4.41(m,1H), 5.15(m), 7.08-7.51(m)

The following compounds were prepared from the appropriate starting materials by the processes described in Example 6.

Example 13

5-t-Butyl-3-[N-(1-(S)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(R)-carboxylic acid

mp 67-9°

Found: C,48.37; H,7.99; N,9.42; S,7.2%

C₂₂H₃₂N₂O₅S · 2H₂O requires

C,48.22; H,7.80; N,9.93; S,7.57%

Example 14

5-t-Butyl-3-[N-(1-(R)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(R)-carboxylic acid

mp 67-9°

Found: C,51.44; H,7.52; N,10.38; S,7.79%

C₂₂H₃₂N₂O₅S · 0.5H₂O requires

C,51.52; H,7.58; N,10.61; S,8.08%

Example 15

5-t-Butyl-3-[N-(1-(R)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid

mp 124-5°

Found: C,52.46; H,7.63; N,10.77; S,8.07%

C₂₂H₃₂N₂O₅S requires

C,52.71; H,7.49; N,10.85; S,8.27%

The following compounds were prepared by the method of Example 5 (using appropriate starting materials)

Example 16

3-[N-(1-(S)-Carboxy-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-(S)-carboxylic acid

mp softens at 151°, decomposes at 165-170°

A fast atom bombardment mass spectrum showed M⁺442 (base peak 91)

5

C₂₂H₂₂N₂O₅S requires MWt 441

Example 17

10

5-t-Butyl-3-[N-(1-(S)-carboxy-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid

15

mp softens at 161°, decomposes at 179-184°

A fast atom bombardment mass spectrum showed M⁺442 (base peak 91)

20

C₂₂H₂₇N₂O₅S requires MWt 421

Example 18

25

5-t-Butyl-3-[N-(1-(S)-carboxybutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid

mp 156-9°

30

The following compounds were prepared by the method of Example 4 (using appropriate starting materials).

35

Example 19

5-Cyclohexyl-3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid

40

mp 136-138°

Example 20

45

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-(pyridin-3-yl)-1,3,4-thiadiazole-2-(S)-carboxylic acid

50

mp 160-3° (softens at c. 140°).

Example 21

55

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2, 3-dihydro-5-isopropyl-1,3,4-thiadiazole-2-(S)-carboxylic acid

Example 22

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-methyl-1,3,4-thiadiazole-2-(S)-carboxylic acid

mp 161-2°

Example 23

5-t-Butyl-3-[N²-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-lysyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid hydrochloride

A solution of 5-t-butyl-3-[N²-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-lysyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid - (0.97g, prepared using the appropriate starting material by the process of Example 4) in tetrahydrofuran (20ml) and water (20ml) was treated with 1N hydrochloric acid (3.8ml). The solvents were evaporated and the residue was taken up in a mixture of dichloromethane and toluene. The solvents were removed by evaporation to give the title product (0.9g) as a white solid.

A mass spectrum (FAB) showed M⁺ 507 (base peak 84).

C₂₃H₃₃N₃O₅S requires MWt 506.

Example 24

5-t-Butyl-3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(R)-carboxylic acid

Prepared from the product of Example 4 step b) by the process of Example 4 step d).

mp 60-63°

Example 25

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-(morpholin-4-yl)-1,3,4-thiadiazole-2-(S)-carboxylic acid maleic acid salt

A solution of 3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-(morpholin-4-yl)-1,3,4-thiadiazole-2-(S)-carboxylic acid (0.47g) - (prepared from the appropriate starting materials by the process of Example 4) in ethanol (25ml) was treated with maleic acid (0.11g). The solvent was removed by evaporation. Trituration of the residue with acetonitrile gave the title product as a white solid (0.31g). mp 174-175°

The following compounds were prepared by the method of Example 3 using appropriate starting materials

Example 26

5-t-Butyl-2,3-dihydro-3-(3-mercapto-1-oxopropyl)-1,3,4-thiadiazole-2-carboxylic acid

mp 128°

Example 27

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-(4-methoxyphenyl)-1,3,4-thiadiazole-2-carboxylic acid

mp 164°

Example 28

Ethyl 3-(3-acetylthio-1-oxopropyl)-2,3-dihydro-5-methylamino-1,3,4-thiadiazole-2-carboxylate

mp 102-103°.

A mass spectrum showed M⁺ 319 (base peak 116).

C₁₇H₁₇N₃O₃S₂ requires MWt 319.

Example 29

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-(2-methylphenyl)-1,3,4-thiadiazole-2-carboxylic acid

mp 118-120°.

Example 30

5-(Furan-2-yl)-2,3-dihydro-3-(3-mercapto-1-oxopropyl)-1,3,4-thiadiazole-2-carboxylic acid

mp 105-108°.

Example 31

Ethyl 3-(3-acetylthio-1-oxopropyl)-5-(4-chlorophenyl)-2,3-dihydro-1,3,4-thiadiazole-2-carboxylate

Prepared by similar processes to those of Example 3 steps a) and b). The product was isolated as a clear gum.

NMR spectrum (CDCl_3) of the compound showed characteristic peaks at δ 7.50 (4H,q,aromatic CH_s), δ 2.34 (3H,s,- SCOCH_3) and δ 6.30 (1H,s, heterocyclic CH).

Example 32

Benzyl 3-(3-acetylthio-1-oxopropyl)-5-benzyl-2,3-dihydro-1,3,4-thiadiazole-2-carboxylate

Prepared using appropriate starting materials by process of Example 3 steps a) and b). The product was isolated as an oil.

NMR spectrum (CDCl_3) showed a characteristic signal at δ 6.17 (1H,s,heterocyclic CH).

Example 33

Benzyl 3-(3-acetylthio-1-oxopropyl)-2,3-dihydro-5-(2-phenylethyl)-1,3,4-thiadiazole-2-carboxylate

Prepared by similar processes to those of Example 3 steps a) and b). The product was isolated as an oil.

NMR spectrum (CDCl_3) showed a characteristic signal at δ 6.10 (1H,s,heterocyclic CH).

Example 34

Ethyl 3-(3-acetylthio-1-oxopropyl)-2,3-dihydro-5-(naphthalen-2-yl)-1,3,4-thiadiazole-2-carboxylate

a) Naphthalene-2-carbothioic acid hydrazide

The sub-title product was prepared from appropriate starting materials by the processes of Example 7, steps a), b), c) and d). mp 166-167°

b) Ethyl 2,3-dihydro-5-(naphthalen-2-yl)-1,3,4-thiadiazole-2-carboxylate

Prepared from the product of step a) and ethyl glyoxylate by the process of Example 3, step a). The crude product was used without further purification.

c) Ethyl 3-(3-acetylthio-1-oxopropyl)-2,3-dihydro-5-(naphthalen-2-yl)-1,3,4-thiadiazole-2-carboxylate

Prepared from the crude product of step b) and 3-acetylthiopropionyl chloride by the process of Example 3, step b). mp 107-108°

Mass spectrum (FAB) showed M^+ 417 (base peak 213).

$\text{C}_{20}\text{H}_{20}\text{N}_2\text{O}_2\text{S}_2$ requires MWt 416.

Example 35

5-(Adamant-1-yl)-2,3-dihydro-3-(3-mercapto-1-oxopropyl)-1,3,4-thiadiazole-2-carboxylic acid

a) Methyl 1-adamantanecarbodithioate

A mixture of 1-adamantanecarboxylic acid chloride (9.0g) and 2,4-bis-methylthio-1,2,3,4-dithiaphosphetan-2,4-disulphide (12.9g) in dry benzene was heated under reflux for 5 hours. The solvent was evaporated and the residue purified by flash chromatography to give the sub-title product (6.2g) as a yellow solid.

mp 64.5-66°

b) Adamantane-1-carbothioic acid hydrazide

A solution of the product of step a) (1g) in methanol (50ml) was treated with hydrazine hydrate (0.3g) and the mixture stirred at room temperature for 1 hour. The solvent was evaporated, the residue triturated with water, and the pH adjusted to 7 to give the sub-title product (0.8g) as a white solid.

mp 204-206°

c) Ethyl 5-(adamant-1-yl)-2,3-dihydro-1,3,4-thiadiazole-2-carboxylate

The product of step b) was treated with ethyl glyoxylate by the process of Example 3, step a) to give the sub-title product (1.5g) as an oil.

d) Ethyl 5-(adamant-1-yl)-2,3-dihydro-3-(3-mercapto-1-oxopropyl)-1,3,4-thiadiazole-2-carboxylate

The crude product of step c) was treated with 3-acetylthiopropionyl chloride by the process of Example 3, step b) to give the sub-title product as an oil.

Mass spectrum (FAB) showed M^+ 425 (base peak 221).

$\text{C}_{20}\text{H}_{20}\text{N}_2\text{O}_2\text{S}_2$ required MWt 424.

e) 5-(Adamant-1-yl)-2,3-dihydro-3-(3-mercapto-1-oxopropyl)-1,3,4-thiadiazole-2-carboxylic acid

The product of step d) was treated with potassium hydroxide by the process of Example 3, step c) to give the title product as white solid. mp 183-184°

Mass spectrum (FAB) showed M⁺355 (base peak 221).

C₁₈H₂₂N₂O₃S₂ requires MWt 354.

Example 36

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-methyl-1,3,4-thiadiazole-2-carboxylic acid dicyclohexylamine salt

A solution of dicyclohexylamine (0.5ml) in ether (10ml) was added to a solution of 2,3-dihydro-3-(3-mercapto-1-oxopropyl)-5-methyl-1,3,4-thiadiazole-2-carboxylic acid (0.5g) (prepared from appropriate starting materials by the method of Example 3) in ether (20ml). The solvent was removed by evaporation.

Trituration of the residue with ether gave the title compound as a white solid (0.5g). mp 150-153°

The following compounds were prepared by the method of Example 36 using appropriate starting materials

Example 37

5-Cyclohexyl-2,3-dihydro-3-(3-mercapto-1-oxopropyl)-1,3,4-thiadiazole-2-carboxylic acid dicyclohexylamine salt

mp 174-6°

Example 38

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-methylthio-1,3,4-thiadiazole-2-carboxylic acid dicyclohexylamine salt

mp 150-3°

Example 39

Benzyl 3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-methylthio-1,3,4-thiadiazole-2-(S)-carboxylate

Prepared by similar processes to those of Example 1 from appropriate starting materials. The product was isolated as a clear gum.

The nmr spectrum of the compound showed characteristic signals at delta 2.55 (3H,s,-SCH₃) and 6.32 (1H,s, heterocyclic CH).

Example A

In vitro Assay of inhibitors of Angiotensin Converting Enzyme

The method is based upon that of Cushman and Cheung (1971) but uses a radioactive substrate [glycine-1-¹⁴C] hippuryl-L-histidyl-L-leucine (HHL) whose hydrolysis may be determined by liquid scintillation counting of released [¹⁴C]-hippuric acid. Hydrolysis of 2mM HHL by an extract of rabbit lung acetone powder (Sigma) over a 30 min. incubation period at 37° is followed by acidification of the reaction mixture and extraction of [¹⁴C]-hippurate with ethyl acetate.

Potential inhibitors are tested initially at 0.01mM and if found are retested at lower concentrations to determine an IC₅₀. Dimethyl sulfoxide at 1% final concentration may be used as a solubility aid without affecting enzyme activity. Compounds of special interest are studied at a range of substrate and inhibitor concentrations to determine the type of inhibition and are also tested against other enzymes, eg carboxypeptidase A to establish their specificity for ACE.

Example B

Antihypertensive effects were investigated in conscious spontaneously hypertensive rats (SHR) of the Okamoto strain. Systolic blood pressure and heart rate were measured by the tail cuff method using an electrophygomanometer 1 hour before and 1, 3, 5 and 24 hours after oral dosing with the compound (dose range 0.1 -100 mg/kg p.o.). Percentage changes in each parameter were measured with respect to pretreatment control values.

Example C

	<u>% w/w</u>	<u>Range % w/w</u>
Compound of formula I	5	1-20
Microcrystalline cellulose	50	10-80
Spray dried lactose	37.75	10-80
Magnesium stearate	1	0.25-2
Colloidal silicon dioxide	0.25	0.1-1
Cross linked sodium carboxy methyl cellulose	3	1-5
Hydroxypropylmethylcellulose (coating)	3	1-5

25 Example D

This formulation is made up as a direct compression tablet, or without compression or coating, may be filled into a gelatine capsule.


30

	<u>% w/w</u>	<u>Range % w/w</u>
Compound of formula I	5	1-20
Microcrystalline cellulose	50	10-80
Lactose	35.75	10-80
Polyvinylpyrrolidone	2	1-5
Magnesium stearate	1	0.25-2
Colloidal silicon dioxide	0.25	0.1-1
Cross linked sodium carboxy methyl cellulose	3	1-5
Hydroxypropyl methyl cellulose (coating)	3	1-5

55

This formulation is made up as a granulate and then compressed into a tablet. Alternatively the granules may be filled into a gelatine capsule.

5



in which Y is S, O or NR₁,

n is 0 or 1,

R₁ is hydrogen or alkyl C 1 to 10,

R₂ is hydrogen, alkyl C 1 to 10, cycloalkyl C3 to 10, CF₃, SR₁₀, a 5 or 6 membered heterocyclic group containing one or more S, O or N atoms, NR₄R₅, phenyl or phenylalkyl C7 to 12, the phenyl, phenylalkyl and heterocyclic groups optionally being fused to a further phenyl group, the heterocyclic group and any phenyl group optionally being substituted by alkyl C 1 to 6, halogen, alkoxy C 1 to 6, nitro, nitrile, CF₃, SR₆, NR₇R₁₁, or hydroxy,

R₄, R₇ and R₁₁, which may be the same or different, are each hydrogen or alkyl C 1 to 10,

R₄ and R₅, which may be the same or different, are each hydrogen, alkyl C 1 to 10 or phenyl,

R₁₀ is alkyl C 1 to 10,

X₁ is S or O, and

D is a chain comprising from 2 -16 atoms, which chain carries an O or S containing substituent at a position 2 -6 atoms away from the group C=X₁,

and pharmaceutically acceptable salts, esters and amides thereof.

2. A compound according to Claim 1, wherein X₁ is O,

D is ZCHR-,

R is hydrogen, alkyl C 1 to 10 or alkyl C 1 to 6 substituted by NH₂,

Z is R₂CH(COOH)NH- or R₂SCH₂-,

Claims

1. A compound of formula I

I

R₁ is hydrogen or R₆CO-,

20 R₆ is alkyl C 1 to 10 or phenyl, and

R₂ is alkyl C 1 to 10 or phenylalkyl C7 to 12.

3. A compound according to Claim 2, wherein Z is R₂CH(COOH)NH- and R₂ is alkyl C 1 to 10 or cycloalkyl C3 to 10.

4. A compound according to Claim 2 wherein

Z is R₂CH(COOH)NH-,

30 Y is S,

R is methyl or aminobutyl,

n is O,

35 R₂ is n-propyl or phenylethyl,

R₂ is t-butyl, and

40 all asymmetric carbon atoms are in the S configuration.

5. 5-t-Butyl-3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

45 5-t-Butyl-3-[N²-(1-(S)-carboxy-3-phenylpropyl)-L-lysyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid, and

50 5-t-Butyl-3-[N-(1-(S)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

and pharmaceutically acceptable salts thereof.

6. 3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-(S)-carboxylic acid,

Benzyl 3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-

L-alanyl]-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-(R)-carboxylate,

Benzyl 3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-(S)-carboxylate,

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-(R)-carboxylic acid,

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-phenyl-1,3,4-thiadiazole-2-carboxylic acid,

Ethyl 3-(3-acetylthio-1-oxopropyl)-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-carboxylate,

5-t-Butyl-3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

Benzyl 5-t-butyl-3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(R)-carboxylate,

Benzyl 5-t-butyl-3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylate,

5-t-Butyl-3-[N²-(1-(S)-carboxy-3-phenylpropyl)-L-lysyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

Benzyl 3-[N⁶-benzyloxycarbonyl-N²-(1-(S)-benzyloxycarbonyl-3-phenylpropyl)-L-lysyl]-5-t-butyl-2,3-dihydro-1,3,4-thiadiazole-2-(R)-carboxylate,

Benzyl 3-[N⁶-benzyloxycarbonyl-N²-(1-(S)-benzyloxycarbonyl-3-phenylpropyl)-L-lysyl]-5-t-butyl-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylate,

5-t-Butyl-3-[N-(1-(S)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

Benzyl 5-t-butyl-3-[N-(1-(S)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(R)-carboxylate,

Benzyl 5-t-butyl-3-[N-(1-(S)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylate,

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-[4-(methylthio)phenyl]-1,3,4-thiadiazole-2-(S)-carboxylic acid,

t-Butyl 3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-[4-(methylthio)phenyl]-1,3,4-thiadiazole-2-(S)-carboxylate,

5 2,3-Dihydro-3-(3-mercapto-2-(S)-methyl-1-oxopropyl)-5-phenyl-1,3,4-thiadiazole-2-(S)-carboxylic acid,

10 Benzyl 3-(3-acetylthio-2-(S)-methyl-1-oxopropyl)-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-carboxylate,

2-Cyclohexyl-5,6-dihydro-4-(3-mercapto-1-oxopropyl)-4H-1,3,4-thiadiazine-5-carboxylic acid,

15 Benzyl 4-(3-acetylthio-1-oxopropyl)-2-cyclohexyl-5,6-dihydro-4H-1,3,4-thiadiazine-5-carboxylate,

20 2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-phenyl-1,3,4-oxadiazole-2-carboxylic acid,

Ethyl 3-(3-acetylthio-1-oxopropyl)-2,3-dihydro-5-phenyl-1,3,4-oxadiazole-2-carboxylate,

25 2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-[4-(trifluoromethyl)phenyl]-1,3,4-thiadiazole-2-carboxylic acid,

30 Benzyl 3-[3-acetylthio-1-oxopropyl]-2,3-dihydro-5-[4-(trifluoromethyl)phenyl]-1,3,4-thiadiazole-2-carboxylate,

Benzyl 4-[3-acetylthio-1-oxopropyl]-5,6-dihydro-1-methyl-2-phenyl-4H-1,3,4-triazine-5-carboxylate,

35 5-t-Butyl-3-[N-(1-(S)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(R)-carboxylic acid,

40 5-t-Butyl-3-[N-(1-(R)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(R)-carboxylic acid,

5-t-Butyl-3-[N-(1-(R)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

45 3-[N-(1-(S)-Carboxy-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-(S)-carboxylic acid,

50 5-t-Butyl-3-[N-(1-(S)-carboxy-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

5-t-Butyl-3-[N-(1-(S)-carboxybutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

55 5-Cyclohexyl-3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-(pyridin-3-yl)-1,3,4-thiadiazole-2-(S)-carboxylic acid,

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-isopropyl-1,3,4-thiadiazole-2-(S)-carboxylic acid,

5-t-Butyl-3-[N²-(1-(S)-ethoxycarbonyl-3-phenyl propyl)-L-lysyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

5-t-Butyl-3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-lysyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-methyl-1,3,4-thiadiazole-2-(S)-carboxylic acid,

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-(morpholin-4-yl)-1,3,4-thiadiazole-2-(S)-carboxylic acid,

5-t-Butyl-2,3-dihydro-3-(3-mercapto-1-oxopropyl)-1,3,4-thiadiazole-2-carboxylic acid,

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-(4-methoxyphenyl)-1,3,4-thiadiazole-2-carboxylic acid,

Ethyl 3-(3-acetylthio-1-oxopropyl)-2,3-dihydro-5-methylamino-1,3,4-thiadiazole-2-carboxylate,

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-(2-methylphenyl)-1,3,4-thiadiazole-2-carboxylic acid,

5-(Furan-2-yl)-2,3-dihydro-3-(3-mercapto-1-oxopropyl)-1,3,4-thiadiazole-2-carboxylic acid,

Ethyl 3-(3-acetylthio-1-oxopropyl)-5-(4-chlorophenyl)-2,3-dihydro-1,3,4-thiadiazole-2-carboxylate,

Benzyl 3-(3-acetylthio-1-oxopropyl)-5-benzyl-2,3-dihydro-1,3,4-thiadiazole-2-carboxylate,

Benzyl 3-(3-acetylthio-1-oxopropyl)-2,3-dihydro-5-(2-phenylethyl)-1,3,4-thiadiazole-2-carboxylate,

Ethyl 3-(3-acetylthio-1-oxopropyl)-2,3-dihydro-5-(naphthalen-2-yl)-1,3,4-thiadiazole-2-carboxylate,

5-(Adamant-1-yl)-2,3-dihydro-3-(3-mercapto-1-oxopropyl)-1,3,4-thiadiazole-2-carboxylic acid,

Ethyl 5-(adamant-1-yl)-2,3-dihydro-3-(3-mercapto-1-oxopropyl)-1,3,4-thiadiazole-2-carboxylate,

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-methyl-1,3,4-thiadiazole-2-carboxylic acid,

5-Cyclohexyl-2,3-dihydro-3-(3-mercapto-1-oxopropyl)-1,3,4-thiadiazole-2-carboxylic acid,

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-methylthio-1,3,4-thiadiazole-2-carboxylic acid, and


Benzyl 3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-methylthio-1,3,4-thiadiazole-2-(S)-carboxylate,

and pharmaceutically acceptable salts thereof.

7. The use of a compound of formula I, as defined in Claim 1, in the production of a pharmaceutical formulation for the treatment of a hypertensive condition.

8. A process for the production of a compound of formula I, as defined in Claim 1, or a pharmaceutically acceptable salt, ester or amide thereof, which comprises


- removal of a protecting group from a compound of formula I in which one or more of the amino or carboxylic acid groups is protected, or
- reaction of a compound of formula II,



II

or a salt, ester, amide, tautomer, or protected derivative thereof,

in which R_3 , Y and n are as defined in Claim 1 with a compound of formula III,




in which D and X, are as defined in Claim 1, and X is a good leaving group,


c) conversion of a compound of formula I, as defined in Claim 1, in which the asymmetric carbon atom of the Y containing heterocyclic ring is in the R configuration into a corresponding compound in which that carbon atom is in the S configuration,

d) reaction of a compound of formula II, in which R_3 , Y and n are as defined in Claim 1,

with a compound of formula VI,



in which D and X, are as defined in Claim 1,



in which Y is S, O or NR_4 ,

n is 0 or 1,

R_4 is hydrogen or alkyl C 1 to 10,

R_3 is hydrogen, alkyl C 1 to 10, cycloalkyl C3 to 10, CF_3 , SR_{10} , a 5 or 6 membered heterocyclic group containing one or more S, O or N atoms, NR_4R_5 , phenyl or phenylalkyl C7 to 12, the phenyl, phenylalkyl and heterocyclic groups optionally being fused to a further phenyl group, the heterocyclic group and any phenyl group optionally being substituted by alkyl C 1 to 6, halogen, alkoxy C 1 to 6, nitro, nitrile, CF_3 , SR_4 , NR_4R_5 , or hydroxy,

R_4 , R_5 and R_{10} , which may be the same or different, are each hydrogen or alkyl C 1 to 10,

e) production of a pharmaceutically acceptable salt of a compound of formula I, as defined in Claim 1, by treating a compound of formula I as defined in Claim 1,

or another salt, an ester or an amide thereof,

with a compound containing an available

pharmaceutically acceptable ion and capable of converting the compound of formula I or the other salt, ester or amide thereof, to a pharmaceutically acceptable salt of the compound of formula I,

and where desired or necessary deprotecting the resulting compound, or converting a compound of formula I to a pharmaceutically acceptable salt, ester or amide thereof or vice versa.

9. A compound of formula II,

II

R_4 and R_5 , which may be the same or different, are each hydrogen, alkyl C 1 to 10 or phenyl,

R_{10} is alkyl C 1 to 10,


and salts, esters, amides and tautomers thereof.

10. A compound according to Claim 1 for use as a pharmaceutical.

11. A pharmaceutical formulation comprising a compound according to Claim 1 in admixture with a pharmaceutically acceptable diluent, excipient or carrier.

Claims for Austria

1. A process for the production of a compound of formula I,



in which Y is S, O or NR₄,

n is 0 or 1,

R₃ is hydrogen or alkyl C 1 to 10,

R₄ is hydrogen, alkyl C 1 to 10, cycloalkyl C3 to 10, CF₃, SR₁₀, a 5 or 6 membered heterocyclic group containing one or more S, O or N atoms, NR₄R₅, phenyl or phenylalkyl C7 to 12, the phenyl, phenylalkyl and heterocyclic groups optionally being fused to a further phenyl group, the heterocyclic group and any phenyl group optionally being substituted by alkyl C 1 to 6, halogen, alkoxy C 1 to 6, nitro, nitrile, CF₃, SR₄, NR₄R₁₁, or hydroxy,

R₄, R₅ and R₁₁, which may be the same or different, are each hydrogen or alkyl C 1 to 10,

I

R₄ and R₅, which may be the same or different, are each hydrogen, alkyl C 1 to 10 or phenyl,


R₁₀ is alkyl C 1 to 10,

X₁ is S or O, and

D is a chain comprising from 2 -16 atoms, which chain carries an O or S containing substituent at a position 2 -6 atoms away from the group C=X₁,

or a pharmaceutically acceptable salt, ester or amide thereof, which comprises,

- a) removal of a protecting group from a compound of formula I in which one or more of the amino or carboxylic acid groups is protected, or
- b) reaction of a compound of formula II,



II

or a salt, ester, amide, tautomer, or protected derivative thereof,

in which R₃, Y and n are as defined in Claim 1 with a compound of formula III,

DC(=X₁)X III

in which D and X₁ are as defined in Claim 1, and X is a good leaving group,

c) conversion of a compound of formula I, as defined in Claim 1, in which the asymmetric carbon atom of the Y containing heterocyclic ring is in the R configuration into a corresponding compound in which that carbon atom is in the S configuration,

d) reaction of a compound of formula II, in which R₃, Y and n are as defined in Claim 1,

with a compound of formula VI,

DC(=X₁)OH VI

in which D and X₁ are as defined in Claim 1,

e) production of a pharmaceutically acceptable salt of a compound of formula I, as defined in Claim 1, by treating a compound of formula I as defined in Claim 1,

or another salt, an ester or an amide thereof,

with a compound containing an available

pharmaceutically acceptable ion and capable of

converting the compound of formula I or the other salt, ester or amide thereof, to a pharmaceutically acceptable salt of the compound of formula I,

and where desired or necessary deprotecting the resulting compound, or converting a compound of formula I to a pharmaceutically acceptable salt, ester or amide thereof or vice versa.

2. A process according to Claim 1, wherein X, is O, D is ZCHR-,

R is hydrogen, alkyl C 1 to 10 or alkyl C 1 to 6 substituted by NH₂,

Z is R₁CH(COOH)NH- or R₁SCH₂-,

R₁ is hydrogen or R₁CO-,

R₂ is alkyl C 1 to 10 or phenyl, and

R₃ is alkyl C 1 to 10 or phenylalkyl C 7 to 12.

3. A process according to Claim 2, wherein Z is R₁CH(COOH)NH- and R₂ is alkyl C 1 to 10 or cycloalkyl C 3 to 10.

4. A process according to Claim 2 wherein

Z is R₁CH(COOH)NH-,

Y is S,

R is methyl or aminobutyl,

n is O,

R₁ is n-propyl or phenylethyl,

R₂ is t-butyl, and

all asymmetric carbon atoms are in the S configuration.

5. A process according to Claim 1, wherein the compound of formula I is,

5-t-Butyl-3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

5-t-Butyl-3-[N²-(1-(S)-carboxy-3-phenylpropyl)-L-lysyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid, and

5-t-Butyl-3-[N-(1-(S)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

and pharmaceutically acceptable salts thereof.

6. A process according to Claim 1, wherein the compound of formula I is,

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-(S)-carboxylic acid,

Benzyl 3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-(R)-carboxylate,

Benzyl 3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-(S)-carboxylate,

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-(R)-carboxylic acid,

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-phenyl-1,3,4-thiadiazole-2-carboxylic acid,

Ethyl 3-(3-acetylthio-1-oxopropyl)-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-carboxylate,

5-t-Butyl-3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

Benzyl 5-t-butyl-3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(R)-carboxylate,

Benzyl 5-t-butyl-3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylate,

5-t-Butyl-3-[N²-(1-(S)-carboxy-3-phenylpropyl)-L-lysyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

Benzyl 3-[N²-benzyloxycarbonyl-N²-(1-(S)-benzyloxycarbonyl-3-phenylpropyl)-L-lysyl]-5-t-butyl-2,3-dihydro-1,3,4-thiadiazole-2-(R)-carboxylate,

Benzyl 3-[N²-benzyloxycarbonyl-N²-(1-(S)-benzyloxycarbonyl-3-phenylpropyl)-L-lysyl]-5-t-butyl-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylate,

5-t-Butyl-3-[N-(1-(S)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

Benzyl 5-t-butyl-3-[N-(1-(S)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(R)-carboxylate,

Benzyl 5-t-butyl-3-[N-(1-(S)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylate,

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-[4-(methylthio)phenyl]-1,3,4-thiadiazole-2-(S)-carboxylic acid,

t-Butyl 3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-[4-(methylthio)phenyl]-1,3,4-thiadiazole-2-(S)-carboxylate,

2,3-Dihydro-3-(3-mercapto-2-(S)-methyl-1-oxopropyl)-5-phenyl-1,3,4-thiadiazole-2-(S)-carboxylic acid,

Benzyl 3-(3-acetylthio-2-(S)-methyl-1-oxopropyl)-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-carboxylate,

2-Cyclohexyl-5,6-dihydro-4-(3-mercapto-1-oxopropyl)-4H-1,3,4-thiadiazine-5-carboxylic acid,

Benzyl 4-(3-acetylthio-1-oxopropyl)-2-cyclohexyl-5,6-dihydro-4H-1,3,4-thiadiazine-5-carboxylate,

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-phenyl-1,3,4-oxadiazole-2-carboxylic acid,

Ethyl 3-(3-acetylthio-1-oxopropyl)-2,3-dihydro-5-phenyl-1,3,4-oxadiazole-2-carboxylate,

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-[4-(trifluoromethyl)phenyl]-1,3,4-thiadiazole-2-carboxylic acid,

Benzyl 3-[3-acetylthio-1-oxopropyl]-2,3-dihydro-5-[4-(trifluoromethyl)phenyl]-1,3,4-thiadiazole-2-carboxylate,

Benzyl 4-[3-acetylthio-1-oxopropyl]-5,6-dihydro-1-methyl-2-phenyl-4H-1,3,4-triazine-5-carboxylate,

5-t-Butyl-3-[N-(1-(S)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(R)-carboxylic acid,

5-t-Butyl-3-[N-(1-(R)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(R)-carboxylic acid,

5-t-Butyl-3-[N-(1-(R)-ethoxycarbonylbutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

3-[N-(1-(S)-Carboxy-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-phenyl-1,3,4-thiadiazole-2-(S)-carboxylic acid,

5-t-Butyl-3-[N-(1-(S)-carboxy-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-

carboxylic acid,

5-t-Butyl-3-[N-(1-(S)-carboxybutyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

5-Cyclohexyl-3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-(pyridin-3-yl)-1,3,4-thiadiazole-2-(S)-carboxylic acid,

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-isopropyl-1,3,4-thiadiazole-2-(S)-carboxylic acid,

5-t-Butyl-3-[N²-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-lysyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

5-t-Butyl-3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-1,3,4-thiadiazole-2-(S)-carboxylic acid,

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-methyl-1,3,4-thiadiazole-2-(S)-carboxylic acid,

3-[N-(1-(S)-Ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-(morpholin-4-yl)-1,3,4-thiadiazole-2-(S)-carboxylic acid,

5-t-Butyl-2,3-dihydro-3-(3-mercapto-1-oxopropyl)-1,3,4-thiadiazole-2-carboxylic acid,

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-(4-methoxyphenyl)-1,3,4-thiadiazole-2-carboxylic acid,

Ethyl 3-(3-acetylthio-1-oxopropyl)-2,3-dihydro-5-methylamino-1,3,4-thiadiazole-2-carboxylate,

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-(2-methylphenyl)-1,3,4-thiadiazole-2-carboxylic acid,

5-(Furan-2-yl)-2,3-dihydro-3-(3-mercapto-1-oxopropyl)-1,3,4-thiadiazole-2-carboxylic acid,

Ethyl 3-(3-acetylthio-1-oxopropyl)-5-(4-chlorophenyl)-2,3-dihydro-1,3,4-thiadiazole-2-carboxylate,

Benzyl 3-(3-acetylthio-1-oxopropyl)-5-benzyl-2,3-dihydro-1,3,4-thiadiazole-2-carboxylate,

Benzyl 3-(3-acetylthio-1-oxopropyl)-2,3-dihydro-5-(2-THIS PARAGRAPH COULD NOT BE READ IN FROM DISK Ethyl 3-(3-acetylthio-1-oxopropyl)-2,3-

dihydro-5-(naphthalen-2-yl)-1,3,4-thiadiazole-2-carboxylate,

5-(Adamant-1-yl)-2,3-dihydro-3-(3-mercapto-1-oxopropyl)-1,3,4-thiadiazole-2-carboxylic acid,

Ethyl 5-(adamant-1-yl)-2,3-dihydro-3-(3-mercapto-1-oxopropyl)-1,3,4-thiadiazole-2-carboxylate,

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-methyl-1,3,4-thiadiazole-2-carboxylic acid,


5-Cyclohexyl-2,3-dihydro-3-(3-mercapto-1-oxopropyl)-1,3,4-thiadiazole-2-carboxylic acid,

2,3-Dihydro-3-(3-mercapto-1-oxopropyl)-5-methylthio-1,3,4-thiadiazole-2-carboxylic acid, and

Benzyl 3-[N-(1-(S)-ethoxycarbonyl-3-phenylpropyl)-L-alanyl]-2,3-dihydro-5-methylthio-1,3,4-thiadiazole-2-(S)-carboxylate,

and pharmaceutically acceptable salts thereof.

7. A compound of formula II,



II

in which Y is S, O or NR₄,

n is 0 or 1,

R₄ is hydrogen or alkyl C 1 to 10,

R₃ is hydrogen, alkyl C 1 to 10, cycloalkyl C3 to 10, CF₃, SR₁₀, a 5 or 6 membered heterocyclic group containing one or more S, O or N atoms, NR₄R₅, phenyl or phenylalkyl C7 to 12, the phenyl, phenylalkyl and heterocyclic groups optionally being fused to a further phenyl group, the heterocyclic group and any phenyl group optionally

being substituted by alkyl C 1 to 6, halogen, alkoxy C 1 to 6, nitro, nitrile, CF₃, SR₄, NR₇R₁₁, or hydroxy,

R₄, R₅ and R₁₁, which may be the same or different, are each hydrogen or alkyl C 1 to 10,

R₄ and R₅, which may be the same or different, are each hydrogen, alkyl C 1 to 10 or phenyl,

R₁₀ is alkyl C 1 to 10,

and salts, esters, amides and tautomers thereof.

EP 86 30 6325

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
A	US-A-4 312 990 (R.D. HAUGWITZ) * Whole document *	1,7-9	C 07 D 285/12 C 07 D 271/10 C 07 D 417/04 A 61 K 31/41

A	EP-A-0 048 159 (UNIVERSITY OF MIAMI) * Claims *	1,7-9	A 61 K 31/535 A 61 K 31/54 A 61 K 37/02 C 07 D 285/16 C 07 D 253/06 C 07 K 5/06 A 61 K 31/53

			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			C 07 D 285/00 C 07 D 271/00 C 07 D 253/00 C 07 D 417/00 C 07 K 5/00
The present search report has been drawn up for all claims			
Place of search		Date of completion of the search	Examiner
THE HAGUE		03-12-1986	ALLARD M.S.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone		T : theory or principle underlying the invention	
Y : particularly relevant if combined with another document of the same category		E : earlier patent document, but published on, or after the filing date	
A : technological background		D : document cited in the application	
O : non-written disclosure		L : document cited for other reasons	
P : intermediate document		& : member of the same patent family, corresponding document	